

STM - structure search
3/13/06

10/795,838

=> d ibib abs hitstr 1-7

L9 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2004:964886 CAPLUS
DOCUMENT NUMBER: 141:417840
TITLE: Transition metal complexes from solid state synthesis
INVENTOR(S): Thummel, Randolph P.; Hu, Yi-Zhen
PATENT ASSIGNEE(S): University of Houston, USA
SOURCE: U.S. Pat. Appl. Publ., 17 pp.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004225122	A1	20041111	US 2004-795838	20040308

PRIORITY APPLN. INFO.: US 2003-453068P P 20030307

AB Methods and comps. related to the synthesis of photosensitizers of **titanium** dioxide performed in situ, stepwise, in the solid state, and directly on the surface of the **titanium** dioxide semiconductor material. The method is generally accomplished by first absorbing an anchor ligand onto the **titanium** dioxide semiconductor material, then adding an appropriate transition metal, and finally adding one or more secondary ligands to complete the synthesis of the transition metal complex or sensitizer. The method is a much simpler and more efficient synthetic technique, than many other preps. In particular, by preparing the sensitizer directly on the surface of the **titanium** dioxide, the lengthy and tedious independent preparation and final absorption of the sensitizer onto the **titanium** dioxide is avoided. Furthermore, the amts. of material required for the in situ solid state synthesis are extremely small; the in situ solid state synthesis method also does not require purification at any stage; unwanted side products or reagents not anchored to the **titanium** dioxide are merely washed away with solvent after each preparative step (Zakeeruddin et al., 1998).

IT 1245-13-2, 4,4'-Dicarboxy-2,2'-biquinoline
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of transition metal complex anchored **titanium** dioxide)

RN 1245-13-2 CAPLUS
CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid (8CI, 9CI) (CA INDEX NAME)



L9 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2003:855556 CAPLUS
DOCUMENT NUMBER: 139:347692
TITLE: Determination of elements in body fluids and test kit including the necessary reagents for diagnosis of

diseaseses caused by elemental imbalances
 INVENTOR(S) : Rupp, Michael E.
 PATENT ASSIGNEE(S) : Future Data, Inc., USA
 SOURCE: U.S. Pat. Appl. Publ., 13 pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2003203495	A1	20031030	US 2003-423130	20030424
US 6821786	B2	20041123		
WO 2003091725	A1	20031106	WO 2003-US12911	20030425
W:				
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,				
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ,				
UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
RW:				
GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				
KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,				
FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,				
BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003223735	A1	20031110	AU 2003-223735	20030425
EP 1504257	A1	20050209	EP 2003-719936	20030425
R:				
AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 2005524071	T2	20050811	JP 2004-500061	20030425
PRIORITY APPLN. INFO.:			US 2002-375566P	P 20020425
			US 2003-423130	A 20030424
			WO 2003-US12911	W 20030425

AB A self-diagnostic test, a self-diagnostic test apparatus, and method of manufacturing

a self-diagnostic test for screening for elemental mineral imbalances in a patient utilizing an anal. of the reaction of mineral specific reagents to a sample from a patient are provided. In one embodiment, the invention is directed to a test for those elements that occur naturally in the body. In such an embodiment, the invention may test for those elements that comprise about 0.001% of the body weight or less (microtrace), those elements that comprise about 4% of the body weight or less (trace), those elements that comprise up to 96% of the body weight (mass), or any combination of the above. A test strip includes series of reagent spots for the colorimetric determination of the individual elements.

IT 979-88-4, Disodium 2,2'-bicinchoninate


RL: ARG (Analytical reagent use); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(determination of elements in body fluids and test kit including the necessary

reagents for diagnosis of diseaseses caused by elemental imbalances)

RN 979-88-4 CAPLUS

CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid, disodium salt (9CI) (CA INDEX NAME)



●2 Na

REFERENCE COUNT: 66 THERE ARE 66 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:23106 CAPLUS

DOCUMENT NUMBER: 138:83329

TITLE: Use of metal ion chelates in validating biological molecules as drug targets in test animal models

INVENTOR(S): Rist, Oystein; Hogberg, Thomas; Holst Lange, Birgitte; Schwartz, Thue W.; Elling, Christian E.

PATENT ASSIGNEE(S): 7TM Pharma A/S, Den.

SOURCE: PCT Int. Appl., 247 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003003009	A1	20030109	WO 2002-DK456	20020628
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, VZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2002054077	A2	20020711	WO 2001-DK867	20011221
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, VZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.:

DK 2001-1026	A	20010629
DK 2001-1027	A	20010629
DK 2001-1028	A	20010629
DK 2001-1030	A	20010629

DK 2001-1031	A	20010629
US 2001-301931P	P	20010629
WO 2001-DK867	A	20011221
WO 2000-EP13389	W	20001229
DK 2001-536	A	20010330
US 2001-280237P	P	20010330

OTHER SOURCE(S): MARPAT 138:83329

AB The invention discloses the use of chemical compds. or selections of chemical compds. (libraries) of the general Formula R1XFY(R1)GZR1 [F, G = N, O, S, Se, P; X, Y, Z = (un)branched C1-12 alkyl, aryl, heteroaryl, etc.; R1 = ABC; A = coupling or connecting moiety; B = spacer moiety; C = functional group] for in vivo methods for testing or validating the physiol. importance and/or the therapeutic or pharmacol. potential of biol. target mols., notably proteins such as, e.g., receptors and especially 7TM receptors

in

test animals expressing the biol. target mol. with, notably, a silent, engineered metal ion site. Use of specific metal ion binding sites of a generic nature in specific biol. target mols. such as, e.g. transmembrane proteins wherein the metal ion binding site is capable of forming a complex with a metal ion is also described. Also disclosed are chemical compds. or libraries suitable for use in methods for improving the in vivo pharmacokinetic behavior of metal ion chelates (e.g. the absorption pattern, the plasma half-life, the distribution, the metabolism and/or the elimination of the metal ion chelates). In order to improve the efficacy of the impact of the metal ion chelate on the biol. target mol. after administration of the metal ion chelate in vivo to a test animal, it is advantageous e.g. to increase the period during which the metal ion chelate is in the circulatory system and/or localized at the target. Further disclosed are metal ion-chelating compds. designed to be suitable for use in a target validation process according to the invention, as well as libraries of at least two or more of such metal ion-chelating compds.


IT 1245-13-2 318512-23-1

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(metal ion chelates in validating biol. mols. as drug targets in test animal models)


RN 1245-13-2 CAPLUS

CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid (8CI, 9CI) (CA INDEX NAME)



RN 318512-23-1 CAPLUS

CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid, di-2-propenyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:23105 CAPLUS

DOCUMENT NUMBER: 138:83328

TITLE: Metal ion binding-based chemical libraries useful for drug discovery processes

INVENTOR(S): Hoegberg, Thomas; Rist, Oystein; Hjelmencrantz, Anders; Moldt, Peter; Elling, Christian E.; Schwartz, Thue W.; Gerlach, Lars Ole; Holst Lange, Birgitte

PATENT ASSIGNEE(S): 7TM Pharma A/S, Den.

SOURCE: PCT Int. Appl., 242 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003003008	A1	20030109	WO 2002-DK455	20020628
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MX, MY, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.:
 DK 2001-1029 A 20010629
 DK 2001-1032 A 20010629
 DK 2001-1033 A 20010629
 DK 2001-1034 A 20010629
 DK 2001-1035 A 20010629
 US 2001-301989P P 20010629
 US 2001-301990P P 20010629

OTHER SOURCE(S): MARPAT 138:83328

AB The invention discloses the use of chemical compds. or selections of chemical compds. (libraries) of the general formula R₁XFY(R₁)GZR₁ [F, G = N, O, S, Se, P; X, Y, Z = (un)branched C₁-12 alkyl, (hetero)aryl, etc.; R₁ = H, ABC; A = coupling or connecting moiety; B = spacer moiety; C = functional group] for in vivo methods for testing or validating the physiol. importance and/or the therapeutic or pharmacol. potential of biol. target mols., notably proteins such as, e.g., receptors and especially 7TM receptors in


test animals expressing the biol. target mol. with, notably, a silent, engineered metal ion site. Use of specific metal ion binding sites of a generic nature in specific biol. target mols. such as, e.g. transmembrane proteins wherein the metal-ion binding site is capable of forming a complex with a metal ion is also described. The invention provides chemical compds. or libraries suitable for use in methods for improving the in vivo pharmacokinetic behavior of metal-ion chelates (e.g. the absorption pattern, the plasma half-life, the distribution, the metabolism and/or the elimination of the metal ion chelates). In order to improve the efficacy of the metal ion chelates impact on the biol. target mol. after administration of the metal ion chelate in vivo to a test animal, it is advantageous e.g. to increase the time period during which the metal ion chelate is in the circulatory system and/or localized at the target. Metal ion chelating compds., which are designed to be suitable for use in a target validation process according to the invention and to libraries of at least two or more of such metal-ion chelating compds. are disclosed.

IT 1245-13-2 318512-23-1

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(metal ion binding-based chemical libraries for drug discovery processes)


RN 1245-13-2 CAPLUS

CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid (8CI, 9CI) (CA INDEX NAME)



RN 318512-23-1 CAPLUS

CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid, di-2-propenyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L9 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2001:839272 CAPLUS

DOCUMENT NUMBER: 136:95093

TITLE: Synthesis and photophysical properties of ruthenium(II) charge transfer sensitizers containing

4,4'-dicarboxy-2,2'-biquinoline and 5,8-dicarboxy-6,7-dihydro-dibenzo[1,10]phenanthroline
 AUTHOR(S): Islam, Ashraful; Sugihara, Hideki; Singh, Lok Pratap; Hara, Kohjiro; Katoh, Ryuzi; Nagawa, Yoshinobu; Yanagida, Masatoshi; Takahashi, Yoshiaki; Murata, Shigeo; Arakawa, Hironori
 CORPORATE SOURCE: National Institute of Materials and Chemical Research (NIMC), Tsukuba, 305-8565, Japan
 SOURCE: Inorganica Chimica Acta (2001), 322(1,2), 7-16
 CODEN: ICHAA3; ISSN: 0020-1693
 PUBLISHER: Elsevier Science S.A.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 136:95093
 AB Ruthenium(II) complexes cis-Ru(H2dcbiq)2X2 and cis-Ru(H2dcdhph)2X2, where H2dcbiq = 4,4'-dicarboxy-2,2'-biquinoline, H2dcdhph = 5,8-dicarboxy-6,7-dihydrodibenzo[1,10]phenanthroline, and X = Cl-, NCS- or CN-, were synthesized and spectroscopically characterized. The resulting complexes show a broad and intense metal-to-ligand charge transfer (MLCT) band in the visible region with a peak between 580 and 700 nm and are emissive at room temperature. The ground-state 1st pKa value of cis-Ru(H2dcbiq)2(NCS)2 (2) is 2.9 by the spectrophotometric method. Photoelectrochem. measurements show that all dyes, when anchored to a nanocryst. TiO2 film electrode, present low light-harvesting efficiencies due to inefficient driving force for electron injection into the conduction band of TiO2 from their lower energy MLCT band. The photoelectrochem. performance of 2 was also studied on a number of oxide semiconductor thin films such as Nb2O5, ZnO, SnO2 and In2O3. The results show that a high value of short-circuit photocurrent (Jsc) is observed for the semiconductors having a low-energy conduction band potential (SnO2 and In2O3). In the dye 2-sensitized TiO2 film, the absorbed photon-to-current conversion efficiency (APCE) spectrum shows an absorption band selective electron injection yield, while a wavelength independent electron injection yield is observed when dye 2 is anchored to SnO2. These results indicate that the lowest excited MLCT state is energetically favorable for electron injection into the conduction band of SnO2 but not for TiO2.
 IT 979-88-4, Disodium 2,2'-biquinoline-4,4'-dicarboxylate
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (reactant; preparation of ruthenium(II) dicarboxybiquinoline and dicarboxydihydrodibenzo[1,10]phenanthroline
 chloro/isothiocyanato/cyanide charge transfer sensitizer complexes)
 RN 979-88-4 CAPLUS
 CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid, disodium salt (9CI) (CA INDEX NAME)



● 2 Na

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1997:753494 CAPLUS

DOCUMENT NUMBER: 128:109926

TITLE: Redox regulation in ruthenium(II) polypyridyl complexes and their application in solar energy conversion

AUTHOR(S): Nazeeruddin, Mohammad K.; Muller, Edgar; Humphry-Baker, Robin; Vlachopoulos, Nick; Gratzel, Michael

CORPORATE SOURCE: Institute of Physical Chemistry, Laboratory for Photonics and Interfaces, Swiss Federal Institute of Technology, Lausanne, CH-1015, Switz.

SOURCE: Journal of the Chemical Society, Dalton Transactions: Inorganic Chemistry (1997), (23), 4571-4578
CODEN: JCDTBI; ISSN: 0300-9246

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Ru(II) complexes [Ru(dmbip)(Hdcbpy)X], [Ru(dmbip)(Hdcbiq)X] and [Ru(dhbip)(Hdcbpy)X], where dmbip = 2,6-bis(1-methylbenzimidazol-2-yl)pyridine, dhbip = 2,6-bis(1-hexadecylbenzimidazol-2-yl)pyridine, H2dcbpy = 4,4'-dicarboxy-2,2'-bipyridine, H2dcbiq = 4,4'-dicarboxy-2,2'-biquinoline and X = Cl⁻, NCS⁻, CN⁻ or H₂O, were synthesized and spectroscopically characterized. They act as efficient charge-transfer sensitizers when anchored onto nanocryst. TiO₂ films. The lowest-energy metal-to-ligand charge-transfer transitions in these complexes could be tuned from 500 to 590 nm by choice of appropriate ligands and the HOMO varied over 400 mV. Some of the complexes reported are emissive at room temperature. The ground- and excited-state pK_a values of dcbpy complexes were measured by spectrophotometric and spectrofluorometric titration. Resonance-Raman spectra show bands characteristic of the dmbip and dcbpy ligand for excitation at 468 nm, while excitation at 568 nm gave predominantly bands associated with the dcbpy ligand. The excited-state pK_a values and the resonance-Raman data indicate that the lowest excited state is a metal to dcbpy or dcbiq ligand charge-transfer state.


IT 1245-13-2, 4,4'-Dicarboxy-2,2'-biquinoline

RL: RCT (Reactant); RACT (Reactant or reagent)

(for preparation of ruthenium(II) polypyridyl complexes as charge-transfer sensitizers on nanocryst. TiO₂ film)

RN 1245-13-2 CAPLUS

CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid (8CI, 9CI) (CA INDEX NAME)



L9 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1997:513073 CAPLUS

DOCUMENT NUMBER: 127:214048

TITLE: Novel sensitizers for photovoltaic cells. Structural variations of Ru(II) complexes containing

10/795,838

AUTHOR(S): 2,6-bis(1-methylbenzimidazol-2-yl)pyridine
Ruille, Stefan; Kohle, Oliver; Pechy, Peter; Graetzel,
Michael
CORPORATE SOURCE: Institut de Chimie Physique, Ecole Polytechnique
Federale de Lausanne, CH-1015, Lausanne, Switz.
SOURCE: Inorganica Chimica Acta (1997), 261(2), 129-140
CODEN: ICHAA3; ISSN: 0020-1693
PUBLISHER: Elsevier
DOCUMENT TYPE: Journal
LANGUAGE: English

AB New Ru(II) complexes were synthesized for the application in photoelectrochem. solar cells based on dye-sensitized nanocryst. TiO₂. Design strategies for the development of new dyes were tested via structural variations of the prototype complex K[Ru(II)(bmipy)(4,4'-dcbpy)(NCS)] (7) with bmipy = 2,6-bis(1-methyl-benzimidazol-2-yl)pyridine, and 4,4'-dcbpy = 2,2'-bipyridine-4,4'-dicarboxylate. A π^* level tuning was performed by replacing 4,4'-dcbpy with the low π^* level ligands dcbiq (2,2'-biquinoline-4,4'-dicarboxylate) and 5,5'-dcbpy. The resulting complexes showed decreased light-to-elec. energy conversion efficiencies. K[Ru(II)(bmipy)(4-PO₃bpy)(NCS)] (4-PO₃H₂bpy = 2,2'-bipyridine-4-phosphonic acid) also sensitized TiO₂ less efficiently than the model compound. The occurrence of two isomers was observed for the complexes containing 4-PO₃bpy.


In

Na[Ru(II)(bmipy)(4,4'-dcbpy)X] with X = substituted phenylcyanamide (pcyd-) anions, the influence of substitution on the Ph group was studied. MLCT absorption maxima of the phenylcyanamide complexes at approx. 510 nm were shifted to lower energies in comparison with the model complex, however photoenergy conversion efficiencies were reduced. When introduced into the complex, phenylcyanamide was coordinated via the nitrile or the amido N. With prolonged reaction time, the amido-bound isomer was partially transformed into the thermodynamically more stable nitrilo-bound isomer. Linkage isomerism of coordinated NCS- and 4-Clpcyd- was studied with multinuclear NMR (1H, 13C, 31P) spectroscopy and 13C-labeled ligands. Prospects for a substantial improvement of Ru(II) polypyridyl sensitizers for TiO₂ are discussed.

IT 1245-13-2, 2,2'-Biquinoline-4,4'-dicarboxylic acid
RL: RCT (Reactant); RACT (Reactant or reagent)
(for preparation of ruthenium benzimidazolylpyridine
biquinolinedicarboxylate complex)

RN 1245-13-2 CAPLUS

CN [2,2'-Biquinoline]-4,4'-dicarboxylic acid (8CI, 9CI) (CA INDEX NAME)



REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d his

(FILE 'HOME' ENTERED AT 12:58:39 ON 13 MAR 2006)

10/795,838

FILE 'REGISTRY' ENTERED AT 12:58:51 ON 13 MAR 2006

L1 STRUCTURE UPLOADED
L2 0 S L1
L3 0 S L1 FULL
L4 STRUCTURE UPLOADED
L5 5 S L4
L6 94 S L4 FULL

FILE 'CAPLUS' ENTERED AT 13:00:36 ON 13 MAR 2006

L7 246 S L6
L8 461803 S TITANIUM?
L9 7 S L7 AND L8

=> d l1
L1 HAS NO ANSWERS
L1 STR

